

Test Form 5

Instructions: You should have with you several number two pencils, an eraser, your 3" x 5" note card, a calculator, and your University ID Card. If you have notes with you, place them in a sealed backpack and place the backpack OUT OF SIGHT or place the notes directly on the table at the front of the room.

Fill in the front page of the Scantron answer sheet with your test form number (listed above), last name, first name, middle initial, and student identification number. **Enter the test form number on your Scantron form, but leave the class section number blank.**

This exam consists of 25 multiple-choice questions. Each question has four points associated with it. Select the best multiple-choice answer by filling in the corresponding circle on the rear page of the answer sheet. If you have any questions before the exam, please ask. If you have any questions during the exam, please ask the proctor. Open and start this exam when instructed. When finished, place your Scantron form and note card in the appropriate stacks. You may keep the exam packet, so please show your work and mark the answers you selected on it.

Abbreviated Solubility Rules:

Rule 1: All nitrates, group 1A metal salts and ammonium salts are soluble.

Rule 2: All carbonates, hydroxides, phosphates and sulfides are insoluble.

Rule 3: Rule 1 always takes precedent.

$R = 0.0821 \frac{L \cdot atm}{mol \cdot K}$	$R = 8.314 \frac{kg \cdot m^2}{s^2 \cdot mol \cdot K}$	$\mu_{rms} = \sqrt{\frac{3RT}{Molar\ Mass}}$
$PV = nRT$	$760\ Torr = 1\ atm = 760\ mm\ Hg$	$K = 273.15 + ^\circ C$
$N_A = 6.02 \times 10^{23}$	$\frac{P_1 V_1}{n_1 T_1} = \frac{P_2 V_2}{n_2 T_2}$	$milli\ (m) = 1/1000$

Substance	J/g · °C
Water	4.184
Methyl Alcohol	2.549
Ice	2.093
Steam	2.009
Benzene	1.750
Wood (typical)	1.674

Substance	J/g · °C
Soil (typical)	1.046
Air	1.046
Aluminum	0.901
Mercury	0.138
Gold	0.130
Lead	0.128

1. A student calculates that 120.04 grams of carbon dioxide should theoretically be produced from the combustion of propane during a process. She actually recovers 112.5 grams of carbon dioxide. What is the percent yield for this process?

- (A) 7.540 %
 (B) 6.281 %
 (C) 6.700 %
 (D) 93.72 %
 (E) 17.54 %

$$\text{Percent Yield} = \left(\frac{\text{Actual}}{\text{Theoretical}} \right) (100\%)$$

$$= \left(\frac{112.5 \text{ g}}{120.04 \text{ g}} \right) (100\%) = 93.72\%$$

2. What is the mass percent composition of ethanol, C₂H₆O?

- (A) %C = 52.14%; %H = 13.13%; %O = 34.73%
 (B) %C = 33.33%; %H = 33.33%; %O = 33.33%
 (C) %C = 22.22%; %H = 66.67%; %O = 11.11%
 (D) %C = 2.22%; %H = 6.67%; %O = 1.11%
 (E) %C = 26.07%; %H = 2.18%; %O = 69.46%


$$\begin{aligned} \text{C} &= 2 \times 12.011 \text{ g/mol} = 24.022 \text{ g/mol} \\ \text{H} &= 6 \times 1.0079 \text{ g/mol} = 6.0474 \text{ g/mol} \\ \text{O} &= 1 \times 16.00 \text{ g/mol} = 16.00 \text{ g/mol} \\ \hline &46.07 \text{ g/mol} \end{aligned}$$

$$\% \text{C} = \frac{24.022 \text{ g/mol}}{46.07 \text{ g/mol}} = 52.14\%$$

$$\% \text{H} = \frac{6.0474 \text{ g/mol}}{46.07 \text{ g/mol}} = 13.13\%$$

$$\% \text{O} = \frac{16.00 \text{ g/mol}}{46.07 \text{ g/mol}} = 34.73\%$$

3. How many grams of H₂O (g) are produced from 440.96 g of propane and excess oxygen?



$$\textcircled{1} \quad 440.96 \text{ g} \left(\frac{1 \text{ mol}}{44.096 \text{ g}} \right) = 10.000 \text{ mol C}_3\text{H}_8$$

$$\textcircled{2} \quad 10.000 \text{ mol C}_3\text{H}_8 \left(\frac{4 \text{ mol H}_2\text{O}}{1 \text{ mol C}_3\text{H}_8} \right) = 40.000 \text{ mol H}_2\text{O}$$


$$\textcircled{3} \quad 40.000 \text{ mol H}_2\text{O} \left(\frac{18.016 \text{ g}}{1 \text{ mol}} \right) = 720.63 \text{ g H}_2\text{O}$$

- (A) 1763.8 g H₂O (g) are produced.
 (B) 44.096 g H₂O (g) are produced.
 (C) 720.63 g H₂O (g) are produced.
 (D) 10.000 g H₂O (g) are produced.
 (E) 180.16 H₂O (g) are produced.

8. A student mixes two solutions: $K_3PO_4(aq)$ and $Ca(NO_3)_2(aq)$. The solid precipitate formed is:


- (A) $KNO_3(s)$
 (B) $Ca_3(PO_4)_2(s)$
 (C) $KOH(s)$
 (D) $CaO(s)$
 (E) $K_3PO_4(s)$

Net ionic equation:



From the solubility rules, PO_4^{3-} is insoluble. It was introduced with a soluble group 1 metal (K^+). It precipitated with Ca^{2+} .

9. A student obtains 25.00 mL of an HCl solution of unknown concentration. Upon titration, 12.74 mL of 0.09950 M NaOH are required for neutralization. Determine the concentration of the HCl solution.



$$M_{HCl} V_{HCl} = M_{NaOH} V_{NaOH}$$

$$(M_{HCl})(25.00 \text{ mL}) = (0.09950 \text{ M})(12.74 \text{ mL})$$

$$M_{HCl} = 0.05071 \text{ M}$$

- (A) 18.74 M.
 (B) 0.07459 M.
 (C) 13.41 M.
 (D) 0.1327 M.
 (E) 0.05071 M.

10. A student obtains a 2.00 liter balloon at 30.0°C . He cools the balloon to -20.0°C . The volume of the balloon at -20.0°C is:

- (A) 1.00 L.
 (B) -2.00 L.
 (C) 1.67 L.
 (D) 2.32 L.
 (E) 1.73 L.

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

Balloon - constant P
 Closed container - constant n

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \quad \frac{2.00 \text{ L}}{303.15 \text{ K}} = \frac{V_2}{253.15 \text{ K}}$$

$$V_2 = 1.67 \text{ L}$$

11. A student places 64.00 grams of oxygen gas (O_2) into a 3.000-L flask at 293.15 K. The pressure inside the flask is:


- (A) 513.4 atm.
 (B) 35.03 atm.
 (C) 16.05 atm.
 (D) 7.978 atm.
 (E) 0.2493 atm.

$$PV = nRT$$

$$P = \frac{nRT}{V} = \frac{\left(\frac{64.00 \text{ g}}{32.00 \text{ g/mol}}\right) \left(0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}}\right) (293.15 \text{ K})}{3.000 \text{ L}}$$

$$P = 16.05 \text{ atm}$$

12. The reaction below takes place in a classroom (a constant pressure of 1.00 atm and a constant temperature of 273 K for both reactants and products).



When 2.00 L of $\text{C}_3\text{H}_8(\text{g})$ react,

- (A) 2.00 L of CO_2 are formed.
 (B) 3.00 L of CO_2 are formed.
 (C) 4.00 L of CO_2 are formed.
 (D) 5.00 L of CO_2 are formed.
 (E) 6.00 L of CO_2 are formed.
13. A student places 3.388 g of a noble gas into a 2.00-L container at 293 K and measures the pressure to be 0.486 atm. This noble gas is:
- (A) He. $PV = nRT \quad n = \frac{PV}{RT} = \frac{(0.486 \text{ atm})(2.00 \text{ L})}{(0.0821 \frac{\text{L}\cdot\text{atm}}{\text{mol}\cdot\text{K}})(293 \text{ K})} = 0.04041 \text{ mol}$
 (B) Ne.
 (C) Ar.
 (D) Kr. $\text{Molar Mass} = \frac{g}{\text{mol}} = \frac{3.388 \text{ g}}{0.04041 \text{ mol}} = 83.8 \text{ g/mol}$
 (E) Xe.
- 36
 Kr
 83.80
14. Consider a sealed balloon containing nitrogen gas. Which of the following is **false**?

- (A) When the temperature is increased, the velocity of the gas molecules increases. True
 (B) When the temperature is increased, the volume of the balloon increases. True
 (C) When the temperature is increased, the moles of gas inside the balloon increases. False
 (D) A 22.4-L balloon, at 1.00 atm, and 273.15 K contains one mole of nitrogen gas. True

15. A student places 2.00 moles of $O_2(g)$ and 4.00 moles of $CH_4(g)$ into a 44.8-L flask at 273 K. The pressure of $CH_4(g)$ is:

- (A) 1/3 atm.
 (B) 1.00 atm.
 (C) 2.00 atm.
 (D) 3.00 atm.
 (E) 2/3 atm.

$$P_{CH_4} = \frac{n_{CH_4} RT}{V} = \frac{(4.00 \text{ mol}) \left(0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \right) (273 \text{ K})}{44.8 \text{ L}}$$

$$P_{CH_4} = 2.00 \text{ atm}$$

16. The root-mean-square speed of $F_2(g)$ at 1.00 atm and 293 K is:

- (A) 13.9 m/s.
 (B) 439 m/s.
 (C) 514 m/s.
 (D) 1191 m/s.
 (E) 192 m/s.

$$u_{rms} = \sqrt{\frac{3RT}{\text{Molar Mass}}} = \sqrt{\frac{(3) \left(8.314 \frac{\text{J}}{\text{mol} \cdot \text{K}} \right) (293 \text{ K})}{38 \times 10^{-3} \frac{\text{kg}}{\text{mol}}}}$$

$$u_{rms} = 439 \frac{\text{m}}{\text{s}}$$

17. Consider the following five gases: $H_2(g)$ $CO_2(g)$ $Ar(g)$ $SF_6(g)$ $Cl_2(g)$

Of these, the gas molecule with the **greatest** velocity at room temperature is:

- (A) $H_2(g)$.
 (B) $CO_2(g)$.
 (C) $Ar(g)$.
 (D) $SF_6(g)$.
 (E) $Cl_2(g)$.


↳ Lightest (Lowest Mass)

23. The heat of formation (ΔH°_f) of $\text{NH}_4\text{Cl}(\text{s})$ is -315.4 kJ/mol . The chemical equation associated with this reaction is:

- (A) $\text{NH}_4(\text{s}) + \text{Cl}(\text{g}) \rightarrow \text{NH}_4\text{Cl}(\text{s})$
(B) $\frac{1}{2} \text{N}_2(\text{g}) + 2 \text{H}_2(\text{g}) + \frac{1}{2} \text{Cl}_2(\text{g}) \rightarrow \text{NH}_4\text{Cl}(\text{s})$
(C) $\text{NH}_4^+(\text{aq}) + \text{Cl}^-(\text{aq}) \rightarrow \text{NH}_4\text{Cl}(\text{s})$
(D) $\text{NH}_4^+(\text{s}) + \text{Cl}^-(\text{s}) \rightarrow \text{NH}_4\text{Cl}(\text{s})$
(E) $\frac{1}{2} \text{N}_2(\text{g}) + 4 \text{HCl}(\text{aq}) \rightarrow \text{NH}_4\text{Cl}(\text{s}) + 3/2 \text{Cl}_2(\text{g})$



24. When the following reaction is carried out in a flask, the flask feels HOT when held in the hands:



Which of the following is **TRUE**?

- (A) Heat is transferred from the flask to the hand; this is an exothermic reaction.
(B) Heat is transferred from the flask to the hand; this is an endothermic reaction.
(C) Heat is transferred from the hand to the flask; this is an exothermic reaction.
(D) Heat is transferred from the hand to the flask; this is an endothermic reaction.



25. Which one of the following statements is **FALSE**?

- (A) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
(B) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
(C) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
(D) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
(E) ΔH for a process is -322.3 kJ . The process is endothermic.

Hint:

The CH 121 Final Exam is scheduled for Thursday, December 9 from 4:00-5:50pm. Rooms will be assigned and posted near the conclusion of the term. There is no opportunity to reschedule the final exam.