

**Test Form 7**

Instructions: You should have with you several number two pencils, an eraser, your 3" x 5" note card, a calculator, and your University ID Card. If you have notes with you, place them in a sealed backpack and place the backpack OUT OF SIGHT or place the notes directly on the table at the front of the room.

Fill in the front page of the Scantron answer sheet with your test form number (listed above), last name, first name, middle initial, and student identification number. **Enter the test form number on your Scantron form, but leave the class section number blank.**

This exam consists of 25 multiple-choice questions. Each question has four points associated with it. Select the best multiple-choice answer by filling in the corresponding circle on the rear page of the answer sheet. If you have any questions before the exam, please ask. If you have any questions during the exam, please ask the proctor. Open and start this exam when instructed. When finished, place your Scantron form and note card in the appropriate stacks. You may keep the exam packet, so please show your work and mark the answers you selected on it.

**Abbreviated Solubility Rules:**

Rule 1: All nitrates, group 1A metal salts and ammonium salts are soluble.

Rule 2: All carbonates, hydroxides, phosphates and sulfides are insoluble.

Rule 3: Rule 1 always takes precedent.

$R = 0.0821 \frac{L \cdot atm}{mol \cdot K}$	$R = 8.314 \frac{kg \cdot m^2}{s^2 \cdot mol \cdot K}$	$\mu_{rms} = \sqrt{\frac{3RT}{Molar Mass}}$
$PV = nRT$	760 Torr = 1 atm = 760 mm Hg	$K = 273.15 + ^\circ C$
$N_A = 6.02 \times 10^{23}$	$\frac{P_1V_1}{n_1T_1} = \frac{P_2V_2}{n_2T_2}$	milli (m) = 1/1000

Substance	J/g. °C
Water	4.184
Methyl Alcohol	2.549
Ice	2.093
Steam	2.009
Benzene	1.750
Wood (typical)	1.674

Substance	J/g. °C
Soil (typical)	1.046
Air	1.046
Aluminum	0.901
Mercury	0.138
Gold	0.130
Lead	0.128

# Periodic Table of the Elements

Periods																		Group	Noble Gases
	IA	IIA	IVA	VA	VIA	VIIA	He											VIIA	
1	H Hydrogen 1.0079																	He Helium 4.0026	
2	Li Lithium 6.941	Be Beryllium 9.01218																He Helium 4.0026	
3	Na Sodium 22.98977	Mg Magnesium 24.305	Al Aluminum 26.9815	B Boron 10.81	C Carbon 12.011	N Nitrogen 14.0067	O Oxygen 15.9994	F Fluorine 18.9984	Ne Neon 20.179									He Helium 4.0026	
4	K Potassium 39.0983	Ca Calcium 40.08	Sc Scandium 44.9559	Ti Titanium 47.88	V Vanadium 50.9415	Cr Chromium 51.986	Mn Manganese 54.9380	Fe Iron 55.847	Co Cobalt 58.9352	Ni Nickel 58.70	Cu Copper 63.546	Zn Zinc 65.38	Ga Gallium 69.72	Ge Germanium 72.59	As Arsenic 74.9216	Se Selenium 78.965	Br Bromine 79.904	Kr Krypton 83.80	
5	Rb Rubidium 85.4678	Sr Strontium 87.62	Y Yttrium 88.9059	Zr Zirconium 91.22	Nb Niobium 92.9064	Mo Molybdenum 95.94	Tc Technetium 98.906	Ru Ruthenium 101.07	Pd Rhodium 102.9055	Rh Rhodium 106.4	Pd Rhodium 107.868	Ag Silver 112.41	In Indium 114.82	Sn Tin 118.69	Te Antimony 121.75	I Tellurium 127.60	Xe Xenon 126.9045		
6	Cs Cesium 132.9054	Ba Barium 137.33	*Rare earths	Hf Hafnium 178.49	Ta Tantalum 180.9479	W Tungsten 183.35	Re Rhenium 186.207	Os Osmium 190.2	Ir Iridium 192.22	Pt Platinum 195.09	Ir Iridium 196.9665	Hg Mercury 200.59	Tl Thallium 204.37	Pb Lead 207.2	Bi Bismuth (209)	Po Polonium (210)	At Astatine (210)	Rn Radon (222)	
7	Fm Francium (223)	Ra Radium (226.0264)	*Actinides	Rutherfordium (261)	Hs Seaborgium (263)	Ns Neilsbohrium (262)	Hs Hassium (265)	Mt Meitnerium (266)	(269)									Stable region?	
	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71				
*Lanthanides	La Lanthanum 138.9055	Ce Cerium 140.12	Pr Praseodymium 140.9077	Nd Neodymium 144.24	Pm Promethium 145	Sm Samarium 150.4	Eu Europium 151.96	Gd Gadolinium 157.25	Tb Terbium 158.9254	Dy Dysprosium 162.50	Ho Holmium 164.9304	Er Erbium 167.26	Tm Thulium 168.9342	Yb Ytterbium 172.04	Lu Lutetium 174.967				
AC	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103				
	Ac Actinium 227.0273	Th Thorium 232.0381	Pa Protactinium 231.0359	U Uranium 238.0299	Np Neptunium 237.0462	Pu Plutonium (244)	Am Americium (243)	Cm Curium (247)	Bk Berkelium (251)	Cf Californium (254)	Es Einsteinium (257)	Fm Fermium (258)	Md Mendelevium (259)	No Nobelium 259	Lr Lawrencium 262				

Note: The atomic mass value given is for naturally occurring proportions of isotopes. Values in parentheses are mass numbers for the most stable isotope.

\*Reported but not confirmed; no name proposed.

1. A student calculates that 120.04 grams of carbon dioxide should theoretically be produced from the combustion of propane during a process. She actually recovers 112.5 grams of carbon dioxide. What is the percent yield for this process?

- (A) 7.540 %.  
 (B) 6.281 %.  
 (C) 6.700 %.  
 (D) 93.72 %.  
 (E) 17.54 %.

$$\text{Percent Yield} = \left( \frac{\text{Actual}}{\text{Theoretical}} \right) \times 100\%$$

$$= \left( \frac{112.5}{120.04} \right) \times 100\% = 93.72\%$$

2. What is the mass percent composition of ethanol, C<sub>2</sub>H<sub>6</sub>O?

- (A) %C = 52.14%;    %H = 13.13%;    %O = 34.73%  
 (B) %C = 33.33%;    %H = 33.33%;    %O = 33.33%  
 (C) %C = 22.22%;    %H = 66.67%;    %O = 11.11%  
 (D) %C = 2.22%;    %H = 6.67%;    %O = 1.11%  
 (E) %C = 26.07%;    %H = 2.18%;    %O = 69.46%


$$\begin{aligned} C &= 2 \times 12.01 \text{ g/mol} = 24.022 \text{ g/mol} \\ H &= 6 \times 1.0079 \text{ g/mol} = 6.0474 \text{ g/mol} \\ O &= 1 \times 16.00 \text{ g/mol} = 16.00 \text{ g/mol} \\ &\hline 46.07 \text{ g/mol} \end{aligned}$$

$$\% C = \frac{24.022 \text{ g/mol}}{46.07 \text{ g/mol}} = 52.14\%$$

$$\% H = \frac{6.0474 \text{ g/mol}}{46.07 \text{ g/mol}} = 13.13\%$$

$$\% O = \frac{16.00 \text{ g/mol}}{46.07 \text{ g/mol}} = 34.73\%$$

3. How many grams of H<sub>2</sub>O (g) are produced from 440.96 g of propane and excess oxygen?



$$\textcircled{1} \quad 440.96 \text{ g} \left( \frac{1 \text{ mol}}{44.096 \text{ g}} \right) = \underline{10.000 \text{ mol C}_3\text{H}_8}$$

$$\textcircled{2} \quad 10.000 \text{ mol C}_3\text{H}_8 \left( \frac{4 \text{ mol H}_2\text{O}}{1 \text{ mol C}_3\text{H}_8} \right) = \underline{40.000 \text{ mol H}_2\text{O}}$$

$$\textcircled{3} \quad 40.000 \text{ mol H}_2\text{O} \left( \frac{18.016 \text{ g}}{1 \text{ mol}} \right) = \underline{720.63 \text{ g H}_2\text{O}}$$

- (A) 1763.8 g H<sub>2</sub>O (g) are produced.  
 (B) 44.096 g H<sub>2</sub>O (g) are produced.  
 (C) 720.63 g H<sub>2</sub>O (g) are produced.  
 (D) 10.000 g H<sub>2</sub>O (g) are produced.  
 (E) 180.16 H<sub>2</sub>O (g) are produced.


4. Which of the following selections contains only acids?

- (A) CH<sub>4</sub>, CH<sub>3</sub>CH<sub>3</sub>, CH<sub>3</sub>CH<sub>2</sub>CH<sub>3</sub>, CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>. Alkanes - not acids or bases  
(B) HNO<sub>3</sub>, NaNO<sub>3</sub>, HCl, NaCl. Acid / salt / Acid / salt  
(C) NaOH, KOH, NH<sub>4</sub>OH, Ca(OH)<sub>2</sub>. Bases  
(D) H<sub>2</sub>SO<sub>4</sub>, HNO<sub>3</sub>, HCl, CH<sub>3</sub>COOH. Strong and weak acids  
(E) HNO<sub>3</sub>, HCl, NH<sub>3</sub>.  
    ↑ Base

5. Consider fuel cells. Which of the following is false?

- (A) The fuel cell consists of tiny chambers that allow hydrogen gas to explode. False.  
(B) The hydrogen fuel cell demonstrated in class produced water. True  
(C) The hydrogen fuel cell demonstrated in class contains platinum to facilitate the process. True  
(D) A hydrogen fuel cell produces energy. True  
(E) The hydrogen fuel cell demonstrated in class used hydrogen and oxygen gases. True

6. Consider the following compound:



carboxylic acid group  
(weak acid)

- (A) a strong base.  
(B) a weak base.  
(C) a strong acid.  
(D) a weak acid.

7. A student places 116.9 grams of NaCl (s) into a 1.000-L volumetric flask and then fills to the mark with water. This is Solution #1. The student then dilutes 0.5000 liters of Solution #1 to a total volume of 1.000 liter. This is Solution #2.

- (A) The concentration of Solution #1 is 2.000 M; the concentration of Solution #2 is 1.000 M.  
(B) The concentration of Solution #1 is 2.000 M; the concentration of Solution #2 is 2.000 M.  
(C) The concentration of Solution #1 is 2.000 M; the concentration of Solution #2 is 4.000 M.  
(D) The concentration of Solution #1 is 116.9 M; the concentration of Solution #2 is 233.8 M.  
(E) The concentration of Solution #1 is 116.9 M; the concentration of Solution #2 is 58.45 M.


Solution #1     $M = \frac{\text{mol}}{\text{L}} = \left( \frac{116.9 \text{ g}}{58.45 \text{ g/mol}} \right) = \underline{2.000 \text{ M}}$

Solution #2     $M_1 V_1 = M_2 V_2$      $(2.000 \text{ M})(0.5000 \text{ L}) = (M_2)(1.000 \text{ L})$   
 $M_2 = \underline{1.000 \text{ M}}$

8. A student mixes two solutions:  $K_3PO_4$  (aq) and  $Ca(NO_3)_2$  (aq). The solid precipitate formed is:


- (A)  $KNO_3$  (s)
- (B)  $Ca_3(PO_4)_2$  (s)
- (C) KOH (s).
- (D) CaO (s).
- (E)  $K_3PO_4$  (s).

Net ionic equation:



From the solubility rules,  $PO_4^{3-}$  is insoluble. It was introduced with a soluble group 1 metal ( $K^+$ ). It precipitated with  $Ca^{2+}$ .

9. A student obtains 25.00 mL of an HCl solution of unknown concentration. Upon titration, 12.74 mL of 0.09950 M NaOH are required for neutralization. Determine the concentration of the HCl solution.



$$M_{HCl} V_{HCl} = M_{NaOH} V_{NaOH}$$

$$(M_{HCl})(25.00 \text{ mL}) = (0.09950 \text{ M})(12.74 \text{ mL})$$

$$M_{HCl} = 0.05071 \text{ M}$$

- (A) 18.74 M.
- (B) 0.07459 M.
- (C) 13.41 M.
- (D) 0.1327 M.
- (E) 0.05071 M.

10. A student obtains a 2.00 liter balloon at 30.0 °C. He cools the balloon to -20.0 °C. The volume of the balloon at -20.0 °C is:

- (A) 1.00 L.
- (B) -2.00 L.
- (C) 1.67 L.
- (D) 2.32 L.
- (E) 1.73 L.

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

Balloon - constant P  
Closed container - constant n

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \quad \frac{2.0 \text{ L}}{303.15 \text{ K}} = \frac{V_2}{253.15 \text{ K}}$$

$$V_2 = 1.67 \text{ L}$$

11. A student places 64.00 grams of oxygen gas ( $O_2$ ) into a 3.000-L flask at 293.15 K. The pressure inside the flask is:


$$PV = nRT$$

- (A) 513.4 atm.
- (B) 35.03 atm.
- (C) 16.05 atm.
- (D) 7.978 atm.
- (E) 0.2493 atm.

$$P = \frac{nRT}{V} = \frac{\left(\frac{64.00 \text{ g}}{32.00 \text{ g/mol}}\right) \times 0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}} \times 293.15 \text{ K}}{3.000 \text{ L}}$$

$$P = 16.05 \text{ atm}$$

12. The reaction below takes place in a classroom (a constant pressure of 1.00 atm and a constant temperature of 273 K for both reactants and products).



When 2.00 L of  $\text{C}_3\text{H}_8(\text{g})$  react,

- (A) 2.00 L of  $\text{CO}_2$  are formed.
  - (B) 3.00 L of  $\text{CO}_2$  are formed.
  - (C) 4.00 L of  $\text{CO}_2$  are formed.
  - (D) 5.00 L of  $\text{CO}_2$  are formed.
  - (E) 6.00 L of  $\text{CO}_2$  are formed.
13. A student places 3.388 g of a noble gas into a 2.00-L container at 293 K and measures the pressure to be 0.486 atm. This noble gas is:

- (A) He.
- (B) Ne.
- (C) Ar.
- (D) Kr.
- (E) Xe.

$$PV = nRT \quad n = \frac{PV}{RT} = \frac{(0.486 \text{ atm})(2.00 \text{ L})}{(0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}})(293 \text{ K})} = 0.04041 \text{ mol}$$

$$\text{Molar Mass} = \frac{g}{\text{mol}} = \frac{3.388 \text{ g}}{0.04041 \text{ mol}} = 83.8 \text{ g/mol}$$

36  
 Kr  
 83.80

14. Consider a sealed balloon containing nitrogen gas. Which of the following is false?

- (A) When the temperature is increased, the velocity of the gas molecules increases. True
- (B) When the temperature is increased, the volume of the balloon increases. True
- (C) When the temperature is increased, the moles of gas inside the balloon increases. False
- (D) A 22.4-L balloon, at 1.00 atm, and 273.15 K contains one mole of nitrogen gas. True

15. A student places 2.00 moles of O<sub>2</sub> (g) and 4.00 moles of CH<sub>4</sub> (g) into a 44.8-L flask at 273 K. The pressure of CH<sub>4</sub> (g) is:

- (A) 1/3 atm.
- (B) 1.00 atm.
- (C) 2.00 atm.
- (D) 3.00 atm.
- (E) 2/3 atm.

$$P_{\text{CH}_4} = \frac{n_{\text{CH}_4} RT}{V} = \frac{(4.00 \text{ mol})(0.0821 \frac{\text{L} \cdot \text{atm}}{\text{mol} \cdot \text{K}})(273 \text{ K})}{44.8 \text{ L}}$$

$$P_{\text{CH}_4} = 2.00 \text{ atm}$$

16. The root-mean-square speed of F<sub>2</sub> (g) at 1.00 atm and 293 K is:

- (A) 13.9 m/s
- (B) 439 m/s.
- (C) 514 m/s.
- (D) 1191 m/s.
- (E) 192 m/s.

$$v_{\text{rms}} = \sqrt{\frac{3RT}{\text{Molar Mass}}} = \sqrt{\frac{(3)(8.314 \frac{\text{J}}{\text{mol} \cdot \text{K}})(293 \text{ K})}{38 \times 10^{-3} \frac{\text{kg}}{\text{mol}}}}$$

$$v_{\text{rms}} = 439 \frac{\text{m}}{\text{s}}$$

17. Consider the following five gases: H<sub>2</sub> (g)      CO<sub>2</sub> (g)      Ar (g)      SF<sub>6</sub> (g)      Cl<sub>2</sub> (g)

Of these, the gas molecule with the greatest velocity at room temperature is:

↳ Lightest (Lowest Mass)


- (A) H<sub>2</sub> (g).
- (B) CO<sub>2</sub> (g).
- (C) Ar (g).
- (D) SF<sub>6</sub> (g).
- (E) Cl<sub>2</sub> (g).

18. Which of the following processes is endothermic?
- (A)  $2 \text{C}_8\text{H}_{18}(\text{l}) + 25 \text{O}_2(\text{g}) \rightarrow 16 \text{CO}_2(\text{g}) + 18 \text{H}_2\text{O}(\text{g})$ . Combustion - exothermic  
 (B)  $\text{NH}_4\text{NO}_3(\text{s}) \rightarrow \text{NH}_4\text{NO}_3(\text{aq})$ . Endothermic - "Cold Pack" reaction  
 (C)  $\text{H}_2\text{O}(\text{g}) \rightarrow \text{H}_2\text{O}(\text{l})$ . Condensation of steam - exothermic  
 (D)  $\text{H}_2\text{O}(\text{l}) \rightarrow \text{H}_2\text{O}(\text{s})$ . Water freezing - exothermic - heat leaves the system
19. How much heat is required to raise the temperature of 2500.0 grams of aluminum from 30.5 °C to 80.0°C?
- $$q = mc\Delta T = (2500.0 \text{ g})(0.901 \frac{\text{J}}{\text{g}\cdot\text{°C}})(80.0^\circ\text{C} - 30.5^\circ\text{C})$$
- $$= \cancel{111498} \quad 111498.75 \text{ J}$$
- $$= 111 \text{ kJ}$$
- (A) 76.3 kJ  
 (B) 517 kJ  
 (C) 111 kJ  
 (D) 8.10 kJ  
 (E) 16.1 kJ
20. A system takes in 40 kJ of heat and does 30 kJ of work. The change in the energy of the system is:
- $$E = q + w$$
- $$= (+40 \text{ kJ}) + (-30 \text{ kJ}) = +10 \text{ kJ}$$
- (A) -70 kJ.  
 (B) +70 kJ.  
 (C) -10 kJ.  
 (D) +10 kJ.  
 (E) 1.33 kJ.
- ↑  
 heat enters  
 the system  
 ↑  
 work  
 "leaves"  
 the system

21. Use the data in the table below to answer the following question:

	$\Delta H_f^\circ$ (kJ/mol)
CO <sub>2</sub> (g)	-393.5
C <sub>3</sub> H <sub>8</sub> (g)	-104.0
H <sub>2</sub> O (l)	-285.9

What is  $\Delta H^\circ_{\text{reaction}}$  for the following reaction?



-104.0


$\Delta H^\circ = \text{products} - \text{reactants}$

(3)(-393.5) + (4)(-285.9)





$$\begin{aligned} &= [(3 \text{ mol CO}_2)(-393.5 \frac{\text{kJ}}{\text{mol CO}_2}) + (4 \text{ mol H}_2\text{O})(-285.9 \frac{\text{kJ}}{\text{mol H}_2\text{O}})] \\ &\quad - [(1 \text{ mol C}_3\text{H}_8)(-104 \frac{\text{kJ}}{\text{mol C}_3\text{H}_8}) + (5 \text{ mol O}_2)(0 \frac{\text{kJ}}{\text{mol O}_2})] \\ &= -2220.1 \text{ kJ} \end{aligned}$$

- (A) -783.4 kJ.
- (B) -2220.1 kJ.
- (C) -2428.1 kJ.
- (D) +2428.1 kJ.
- (E) +575.4 kJ.

22. Determine  $\Delta H^\circ$  for the reaction C(graphite) + 2 H<sub>2</sub>(g) → CH<sub>4</sub>(g), using:



- (A) -105.5 kJ.
- (B) -74.8 kJ.
- (C) -1570 kJ.
- (D) -211.0 kJ.
- (E) +211.0 kJ.




23. The heat of formation ( $\Delta H^\circ_f$ ) of  $\text{NH}_4\text{Cl}$  (s) is  $-315.4 \text{ kJ/mol}$ . The chemical equation associated with this reaction is:

- (A)  $\text{NH}_4(s) + \text{Cl}(g) \rightarrow \text{NH}_4\text{Cl}(s)$
- (B)  $\frac{1}{2} \text{N}_2(g) + 2 \text{H}_2(g) + \frac{1}{2} \text{Cl}_2(g) \rightarrow \text{NH}_4\text{Cl}(s)$
- (C)  $\text{NH}_4^+(aq) + \text{Cl}^-(aq) \rightarrow \text{NH}_4\text{Cl}(s)$
- (D)  $\text{NH}_4^+(s) + \text{Cl}^-(s) \rightarrow \text{NH}_4\text{Cl}(s)$
- (E)  $\frac{1}{2} \text{N}_2(g) + 4 \text{HCl}(aq) \rightarrow \text{NH}_4\text{Cl}(s) + \frac{3}{2} \text{Cl}_2(g)$



24. When the following reaction is carried out in a flask, the flask feels HOT when held in the hands:



Which of the following is **TRUE**?

- (A) Heat is transferred from the flask to the hand; this is an exothermic reaction.
- (B) Heat is transferred from the flask to the hand; this is an endothermic reaction.
- (C) Heat is transferred from the hand to the flask; this is an exothermic reaction.
- (D) Heat is transferred from the hand to the flask; this is an endothermic reaction.



25. Which one of the following statements is **FALSE**?

- (A) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
- (B) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
- (C) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
- (D) The CH 121 Final Exam is scheduled for Thursday, December 9 at 4:00pm.
- (E)  $\Delta H$  for a process is  $-322.3 \text{ kJ}$ . The process is endothermic.

**Hint:**

The CH 121 Final Exam is scheduled for Thursday, December 9 from 4:00-5:50pm. Rooms will be assigned and posted near the conclusion of the term. There is no opportunity to reschedule the final exam.