

**DO NOT OPEN THIS EXAM UNTIL INSTRUCTED.
CALCULATORS ARE NOT TO BE SHARED.**

Test Form 1

Instructions: You should have with you several number two pencils, an eraser, your 3" x 5" note card, a calculator, and your University ID Card. If you have notes with you, place them in a sealed backpack and place the backpack OUT OF SIGHT or place the notes directly on the table at the front of the room.

Fill in the front page of the Scantron answer sheet with your test form number (listed above), last name, first name, middle initial, and student identification number. **Leave the class section number blank.**

This exam consists of 25 multiple-choice questions. Each question has four points associated with it. Select the best multiple-choice answer by filling in the corresponding circle on the rear page of the answer sheet. If you have any questions before the exam, please ask. If you have any questions during the exam, please ask the proctor. Open and start this exam when instructed. When finished, place your Scantron form in the appropriate stack. You may keep the exam packet, so please show your work and mark the answers you selected on it.

IA

1
H
Hydrogen
1.0079


3 Li Lithium 6.941	4 Be Beryllium 9.01218
11 Na Sodium 22.98977	12 Mg Magnesium 24.305

VIIA

2
He
Helium
4.0026

III A	IV A	V A	VII A	VII A	10 Ne
5 B Boron 10.81	6 C Carbon 12.011	7 N Nitrogen 14.0067	8 O Oxygen 15.9994	9 F Fluorine 18.9984	20.179
13 Al Aluminum 26.9815	14 Si Silicon 28.0855	15 P Phosphorus 30.97376	16 S Sulfur 32.06	17 Cl Chlorine 35.453	18 Ar Argon 39.948
31 Ga Gallium 69.72	32 Ge Germanium 72.59	33 As Arsenic 74.9216	34 Se Selenium 78.96	35 Br Bromine 79.904	36 Kr Krypton 83.80
37 Rb Rubidium 85.4678	38 Sr Strontium 87.62	39 Y Yttrium 88.9059	40 Zr Zirconium 91.22	41 Nb Niobium 92.9064	42 Mo Molybdenum 95.94
43 Tc Technetium 98.906	44 Ru Ruthenium 101.07	45 Rh Rhodium 102.9055	46 Pd Palladium 106.4	47 Ag Silver 107.868	48 Cd Cadmium 112.41
49 In Indium 114.82	50 Sn Tin 118.69	51 Sb Antimony 121.75	52 Te Tellurium 127.60	53 I Iodine 126.9045	54 Xe Xenon 131.30
55 Cs Cesium 132.9054	56 Ba Barium 137.33	57-71 *Rare earths	72 Hf Hafnium 178.49	73 Ta Tantalum 180.9479	74 W Tungsten 183.85
75 Re Rhenium 190.2	76 Os Osmium 192.22	77 Ir Iridium 192.22	78 Pt Platinum 195.09	79 Au Gold 196.9665	80 Hg Mercury 200.59
81 Tl Thallium 204.37	82 Pb Lead 207.2	83 Bi Bismuth 208.9804	84 Po Polonium (209)	85 At Astatine (210)	86 Rn Radon (222)
87 Fr Francium (223)	88 Ra Radium 226.0254	89-103 Actinides (261)	104 Rf Rutherfordium (262)	105 Ha Hahnium (263)	106 Sg Seaborgium (262)
107 Ns Neilsbohrium (262)	108 HS Hassium (265)	109 Mt Meitnerium (266)	110 †	111 ‡	
					114
					→ Stable region?

57 La Lanthanum 138.9055	58 Ce Cerium 140.12	59 Pr Praseodymium 140.9077	60 Nd Neodymium 144.24	61 Pm Promethium 145	62 Sm Samarium 150.4	63 Eu Europium 151.96	64 Gd Gadolinium 157.25	65 Tb Terbium 158.9254	66 Dy Dysprosium 162.50	67 Ho Holmium 164.9304	68 Er Erbium 167.26	69 Tm Thulium 168.9342	70 Yb Ytterbium 173.04	71 Lu Lutetium 174.967
89 Ac Actinium 227.0278	90 Th Thorium 232.0381	91 Pa Protactinium 231.0359	92 U Uranium 238.029	93 Np Neptunium 237.0482	94 Pu Plutonium (244)	95 Am Americium (243)	96 Cm Curium (247)	97 Bk Berkelium (247)	98 Cf Californium (251)	99 Es Einsteinium (254)	100 Fm Fermium (257)	101 Md Mendelevium (258)	102 No Nobelium 259	103 Lr Lawrencium 262



Spectrochemical series: $\text{CN}^- > \text{NO}_2^- > \text{en} > \text{NH}_3 > \text{NCS}^- > \text{H}_2\text{O} > \text{F}^- > \text{Cl}^-$


$F = 96,485 \text{ C/mole e}^-$

$N_A = 6.02 \times 10^{23}$

Reduction Half-Reaction	E° , volt
Acidic Solution	
$F_2(g) + 2 e^- \rightarrow 2F^-(aq)$	+2.866
$O_3(g) + 2 H^+(aq) + 2 e^- \rightarrow O_2(g) + H_2O(l)$	+2.075
$S_2O_8^{2-}(aq) + 2 e^- \rightarrow 2 SO_4^{2-}(aq)$	+2.01
$H_2O_2(aq) + 2H^+(aq) + 2 e^- \rightarrow 2 H_2O(l)$	+1.763
$MnO_4^-(aq) + 8H^+(aq) + 5 e^- \rightarrow Mn^{2+}(aq) + 4 H_2O(l)$	+1.51
$PbO_2(s) + 4H^+(aq) + 2 e^- \rightarrow Pb^{2+}(aq) + 2 H_2O(l)$	+1.455
$Cl_2(g) + 2 e^- \rightarrow 2 Cl^-(aq)$	+1.358
$Cr_2O_7^{2-}(aq) + 14 H^+(aq) + 6 e^- \rightarrow 2 Cr^{3+}(aq) + 7 H_2O(l)$	+1.33
$MnO_2(s) + 4H^+(aq) + 2 e^- \rightarrow Mn^{2+}(aq) + 2 H_2O(l)$	+1.23
$O_2(g) + 4H^+(aq) + 4 e^- \rightarrow 2 H_2O(l)$	+1.229
$2 IO_3^-(aq) + 12H^+(aq) + 10 e^- \rightarrow I_2(s) + 6 H_2O(l)$	+1.20
$Br_2(l) + 2 e^- \rightarrow 2 Br^-(aq)$	+1.065
$NO_3^-(aq) + 4H^+(aq) + 3 e^- \rightarrow NO(g) + 2 H_2O(l)$	+0.956
$Ag^+(aq) + e^- \rightarrow Ag(s)$	+0.800
$Fe^{3+}(aq) + e^- \rightarrow Fe^{2+}(aq)$	+0.771
$O_2(g) + 2H^+(aq) + 2 e^- \rightarrow H_2O_2(aq)$	+0.695
$I_2(s) + 2 e^- \rightarrow 2 I^-(aq)$	+0.535
$Cu^{2+}(aq) + 2 e^- \rightarrow Cu(s)$	+0.340
$SO_4^{2-}(aq) + 4H^+(aq) + 2 e^- \rightarrow 2 H_2O(l) + SO_2(g)$	+0.17
$Sn^{4+}(aq) + 2 e^- \rightarrow Sn^{2+}(aq)$	+0.154
$S(s) + 2H^+(aq) + 2 e^- \rightarrow H_2S(g)$	+0.14
$2H^+(aq) + 2 e^- \rightarrow H_2(g)$	0
$Pb^{2+}(aq) + 2 e^- \rightarrow Pb(s)$	-0.125
$Sn^{2+}(aq) + 2 e^- \rightarrow Sn(s)$	-0.137
$Co^{2+}(aq) + 2 e^- \rightarrow Co(s)$	-0.277
$Fe^{2+}(aq) + 2 e^- \rightarrow Fe(s)$	-0.440
$Zn^{2+}(aq) + 2 e^- \rightarrow Zn(s)$	-0.763
$Al^{3+}(aq) + 3 e^- \rightarrow Al(s)$	-1.676
$Mg^{2+}(aq) + 2 e^- \rightarrow Mg(s)$	-2.356
$Na^+(aq) + e^- \rightarrow Na(s)$	-2.713
$Ca^{2+}(aq) + 2 e^- \rightarrow Ca(s)$	-2.84
$K^+(aq) + e^- \rightarrow K(s)$	-2.924
$Li^+(aq) + e^- \rightarrow Li(s)$	-3.040
Basic Solution	
$O_3(g) + H_2O(l) + 2 e^- \rightarrow O_2(g) + 2 OH^-(aq)$	+1.246
$OCl^-(g) + H_2O(l) + 2 e^- \rightarrow Cl^-(aq) + 2 OH^-(aq)$	+0.890
$O_2(g) + 2 H_2O(l) + 4 e^- \rightarrow 4 OH^-(aq)$	+0.401
$2 H_2O(l) + 2 e^- \rightarrow H_2(g) + 2 OH^-(aq)$	-0.828

1. The oxidation number of each vanadium in $K_2V_2O_6$ is:

- (A) +2
- (B) +3
- (C) +4
- (D) +5
- (E) +6




2. Consider a "General Chemistry Battery" in which one beaker contains aqueous copper sulfate ($CuSO_4$) and a copper metal electrode and the other beaker contains aqueous zinc sulfate ($ZnSO_4$) and a zinc metal electrode.

Which of the following statements is false?

- (A) The mass of the copper electrode will increase as the process proceeds $Cu^{2+}(aq) + 2e^- \rightarrow Cu^0(s)$
- (B) A wire is used to allow the flow of ions but restrict the flow of electrons False
- (C) The concentration of $Zn^{2+}(aq)$ increases as the process proceeds
- (D) Electrons flow from the zinc beaker to the copper beaker $\rightarrow Zn^0(s) \rightarrow Zn^{2+}(aq) + 2e^-$
- (E) The theoretical cell voltage is 1.103 V


\downarrow
Right

Right, Cu^{2+} is higher on the reduction potential table



3. Consider the reaction $Cl_2(g) + 2 Ag(s) \rightarrow 2 Cl^-(aq) + 2 Ag^+(aq)$. The species being oxidized is:


- (A) $Cl_2(g)$
- (B) $Ag(s)$
- (C) $Cl^-(aq)$
- (D) $Ag^+(aq)$



4. Consider fuel cells. Which of the following is false?
- (A) A hydrogen fuel cell produces energy.
 - (B) The hydrogen fuel cell demonstrated in class produced water.
 - (C) The hydrogen fuel cell demonstrated in class contains platinum to facilitate the process.
 - (D) The fuel cell consists of tiny chambers that allow hydrogen gas to explode.
 - (E) The hydrogen fuel cell demonstrated in class input hydrogen and oxygen gases.


5. The calculated cell potential for the $\text{Sn}(\text{s}) + \text{Br}_2(\text{l}) \rightarrow 2 \text{Br}^-(\text{aq}) + \text{Sn}^{2+}(\text{aq})$ cell is:

- (A) + 1.100 V
- (B) + 1.339 V
- (C) + 0.791 V
- (D) + 0.928 V
- (E) + 1.202 V



6. A student provides a current of 4.50 amps through an aqueous solution of $\text{Cu}^{2+}(\text{aq})$ for 2.00 hours. The voltage is such that copper metal is plated. The mass of copper deposited is:


- (A) 7.85 g
- (B) 10.7 g
- (C) 2.73 g
- (D) 12.73 g
- (E) 37.9 g



$$2.00 \times \left(\frac{3600 \text{ s}}{1 \text{ h}} \right) \left(\frac{4.50 \text{ C}}{1 \text{ s}} \right) \left(\frac{1 \text{ mol e}^-}{96,485 \text{ C}} \right) \left(\frac{1 \text{ mol Cu}}{2 \text{ mol e}^-} \right) \left(\frac{63.546 \text{ g}}{1 \text{ mol Cu}} \right) = 10.7 \text{ g Cu}$$




7. Consider Na^+ (aq), Pb^{2+} (aq), Zn^{2+} (aq), Ag^+ (aq), and Li^+ (aq). The strongest oxidizing agent is:

- (A) Na^+ (aq)
- (B) Pb^{2+} (aq)
- (C) Zn^{2+} (aq)
- (D) Ag^+ (aq)
- (E) Li^+ (aq)





8. When the reaction $\text{Fe}(\text{s}) + \text{Cr}_2\text{O}_7^{2-}(\text{aq}) \rightarrow \text{Cr}^{2+}(\text{aq}) + \text{Fe}^{3+}(\text{aq})$ is correctly balanced in acid,

- (A) 3 protons (H^+) are consumed
- (B) 7 protons (H^+) are consumed
- (C) 8 protons (H^+) are consumed
- (D) 12 protons (H^+) are consumed
- (E) 42 protons (H^+) are consumed




9. $^{247}\text{Bk}^{2+}$ has:

- (A) 97 protons, 247 neutrons, 99 electrons
- (B) 150 protons, 97 neutrons, 99 electrons
- (C) 97 protons, 150 neutrons, 95 electrons
- (D) 247 protons, 247 neutrons, 95 electrons
- (E) 247 protons, 238 neutrons, 95 electrons




10. Th-234 decays to produce a beta particle and _____.

- (A) Ra-230
- (B) Th-233
- (C) Th-235
- (D) U-238
- (E) Pa-234




11. Gd-150 decays to produce an alpha particle and _____.

- (A) Gd-146
- (B) Tb-150
- (C) Tb-146
- (D) Sm-150
- (E) Sm-146




12. When a beta particle is emitted,

- (A) An electron is converted to a helium nucleus.
- (B) A gamma ray is released.
- (C) Two gamma rays are released.
- (D) A proton is converted to a neutron.
- (E) A neutron is converted to a proton.



13. A student isolates a sample of tritium ($t_{1/2} = 12.26$ y) containing 5,000 atoms. How much time must pass for only 625 tritium atoms to remain?

- (A) 36.8 y
- (B) 24.5 y
- (C) 12.3 y
- (D) 46.2 y
- (E) 43.6 y



$$\textcircled{1} \quad \ln \frac{1}{2} = -kt_{1/2}$$

$$-0.6931 = -(k)(12.26\text{ y})$$

$$k = 0.0565 \frac{1}{\text{y}}$$

$$\textcircled{2} \quad \ln \frac{A}{A_0} = -kt$$

$$\ln \frac{625 \text{ atoms}}{5000 \text{ atoms}} = -(0.0565 \frac{1}{\text{y}})(t)$$

$$t = 36.8\text{ y}$$





14. Considering the carbon cycle and radiocarbon dating, which of the following statements is false?

- (A) The carbon-14 concentration in fossils is less than the carbon-14 concentration in you.
- (B) Carbon-14 in living organisms does not undergo decay. *It does and it is not replenished*
- (C) Carbon-14 can be used to date specimens previously in the carbon cycle.
- (D) Carbon-14 is generated in the upper atmosphere.

15. Considering nuclear chemistry, which of the following statements is false?


- (A) An example of nuclear fusion is ${}^1_1\text{H} + {}^2_1\text{H} \rightarrow {}^3_2\text{He}$.
- (B) An example of nuclear fission is ${}^1_0\text{n} + {}^{235}_{92}\text{U} \rightarrow {}^{137}_{52}\text{Te} + {}^{97}_{40}\text{Zr} + 2 {}^1_0\text{n}$.
- (C) The half-life is the time required for a sample to decay to one-half its original amount.
- (D) Gamma radiation has a mass of -1.
- (E) A Geiger Counter can be used to show that the orange pigment in certain ceramic glazes is radioactive.

16. Which of the following correctly depicts the curve of nuclear binding energy?




17. Consider the band of stability (AKA "Belt of Stability" located near the beginning of the exam). What decay is expected for a species located to the left of the belt?

- (A) Alpha decay
- (B) Beta decay
- (C) Gamma decay
- (D) Social decay
- (D) UK decay




18. A radioactive decay series that begins with ^{238}U ends with formation of the stable nuclide ^{206}Pb . How many alpha particle emissions and how many beta particle emissions are involved in the sequence of radioactive decays?

- (A) 6 alpha and 8 beta decays
 (B) 6 alpha and 6 beta decays
 (C) 8 alpha and 8 beta decays
 (D) 8 alpha and 6 beta decays
 (E) 8 alpha and 2 beta decays



19. The complex:




- (A) is cis- $[\text{CuCl}_3\text{F}_3]^{4-}$
 (B) is trans- $[\text{CuCl}_3\text{F}_3]^{4-}$
 (C) is fac- $[\text{CuCl}_3\text{F}_3]^{4-}$
 (D) is mer- $[\text{CuCl}_3\text{F}_3]^{4-}$

20. Consider coordination chemistry. Which of the following is false?

- (A) trans- $[\text{FeF}_2\text{Cl}_4]^{3-}$ is nonpolar ✓
 (B) $[\text{CuF}_6]^{3-}$ is a square planar complex Octahedral
 (C) $\text{C}_2\text{O}_4^{2-}$ (ox; the oxalate ion) is a Lewis Base ✓
 (D) $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$ (en; ethylenediamine) is a bidentate ✓
 (E) F^- is a Lewis Base ✓

21. The coordination number for Fe in $[\text{Fe}(\text{ox})_3]^{3-}$ is:

- (A) 1
(B) 2
(C) 3
(D) 4
(E) 6



22. How many d-electrons does Fe^{3+} have?

- (A) 3
(B) 5
(C) 7
(D) 8
(E) 9

$$8 - 3 = 5$$

↑ ↘
group # charge


23. Consider $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$. Which of the following is false?

- (A) $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ is square planar ✓
(B) $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ has mer- and fac- isomers cis- and trans- isomers
(C) cis- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ is polar ✓
(D) The coordination number for Pt in $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ is 4 ✓
(E) trans- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ is non-polar ✓

[Turn over for last page of the exam]

24. How many unpaired electrons are present in $[\text{Mn}(\text{CN})_6]^{4-}$?
[Mn is the Mn^{2+} ion; CN is the CN^- ion; and the Mn^{2+} is low spin].

- (A) 0
(B) 1
(C) 2
(D) 3
(E) 5



25. The CH 123 Final Exam is scheduled for Monday, June 7, 2010, 4:00-5:50pm. Which one of the following statements is FALSE?

- (A) The CH 123 Final Exam is scheduled for Monday, June 7, 2010, 4:00-5:50pm
(B) The CH 123 Final Exam is scheduled for Monday, June 7, 2010, 4:00-5:50pm
(C) The CH 123 Final Exam is scheduled for Monday, June 7, 2010, 4:00-5:50pm
(D) The CH 123 Final Exam is scheduled for Monday, June 7, 2010, 4:00-5:50pm
(E) Your chemistry notes cannot impress your mother enough to entice her into writing a check for \$1000.

[Any response will receive full credit; even no response.]