

## Central Forces Homework 7

Due 5/31/17, 4 pm

**For every problem**, before you start the problem, make a brief statement of the form that a correct solution should have, clearly indicating what quantities you need to solve for. This statement will be graded.

### PRACTICE:

1. Consider the normalized state  $|\Phi\rangle$  for a quantum mechanical particle of mass  $\mu$  constrained to move on a circle of radius  $r_0$ , given by:

$$|\Phi\rangle = \frac{\sqrt{3}}{2} |3\rangle + \frac{i}{2} |-2\rangle$$

- (a) What is the probability that a measurement of  $L_z$  will yield  $2\hbar$ ?  $3\hbar$ ?
  - (b) What is the probability that a measurement of energy will yield  $E = \frac{2\hbar^2}{I}$ ?
  - (c) What is the expectation value of  $L_z$  in this state?
  - (d) What is the expectation value of the energy in this state?
2. Consider the normalized state  $|\Phi\rangle$  for a quantum mechanical particle of mass  $\mu$  constrained to move on a circle of radius  $r_0$ , given by:

$$|\Phi\rangle = \frac{\sqrt{3}}{2} |3\rangle + \frac{i}{2} |-2\rangle$$

- (a) What is the probability that a measurement of  $L_z$  will yield  $2\hbar$ ?  $3\hbar$ ?
- (b) If you measured the z-component of angular momentum to be  $3\hbar$ , what would the state of the particle be immediately after the measurement is made?
- (c) What is the probability that a measurement of energy will yield  $E = \frac{2\hbar^2}{I}$ ?
- (d) What is the expectation value of  $L_z$  in this state?
- (e) What is the expectation value of the energy in this state?
- (f) If you measured the z-component of angular momentum at some time  $t \neq 0$ , what is the probability that you would obtain  $3\hbar$ ?

### REQUIRED:

3. Consider the following normalized quantum state on a ring:

$$\Phi(\phi) = \sqrt{\frac{8}{3\pi}} \sin^2(3\phi) \cos(\phi)$$

- (a) If you measured the  $z$ -component of angular momentum, what is the probability that you would obtain  $\hbar$ ?  $-3\hbar$ ?  $-7\hbar$ ?
  - (b) If you measured the  $z$ -component of angular momentum, what other possible values could you obtain with non-zero probability?
  - (c) If you measured the energy, what is the probability that you would obtain  $\frac{\hbar^2}{2I}$ ?  $\frac{4\hbar^2}{2I}$ ?  $\frac{25\hbar^2}{2I}$ ?
  - (d) If you measured the energy, what possible values could you obtain with non-zero probability?
  - (e) What is the probability that the particle can be found in the region  $0 < \phi < \frac{\pi}{4}$ ? In the region  $\frac{\pi}{4} < \phi < \frac{3\pi}{4}$ ?
  - (f) Plot this wave function.
  - (g) What is the expectation value of  $L_z$  in this state?
4. In this problem, you will carry out calculations on the following normalized abstract quantum state on a ring:

$$|\Psi\rangle = \sqrt{\frac{1}{4}} \left( |1\rangle + \sqrt{2} |2\rangle + |3\rangle \right)$$

- (a) You carry out a measurement to determine the energy of the particle at time  $t = 0$ . Calculate the probability that you measure the energy to be  $\frac{4\hbar^2}{2I}$ .
- (b) You carry out a measurement to determine the  $z$ -component of the angular momentum of the particle at time  $t = 0$ . Calculate the probability that you measure the  $z$ -component of the angular momentum to be  $3\hbar$ .
- (c) You carry out a measurement on the location of the particle at time,  $t = 0$ . Calculate the probability that the particle can be found in the region  $0 < \phi < \frac{\pi}{2}$ .
- (d) You carry out a measurement to determine the energy of the particle at time  $t = \frac{2I}{\hbar} \frac{\pi}{4}$ . Calculate the probability that you measure the energy to be  $\frac{4\hbar^2}{2I}$ .
- (e) You carry out a measurement to determine the  $z$ -component of the angular momentum of the particle at time  $t = \frac{2I}{\hbar} \frac{\pi}{4}$ . Calculate the probability that you measure the  $z$ -component of the angular momentum to be  $3\hbar$ .
- (f) You carry out a measurement on the location of the particle at time,  $t = \frac{2I}{\hbar} \frac{\pi}{4}$ . Calculate the probability that the particle can be found in the region  $0 < \phi < \frac{\pi}{2}$ .
- (g) Write a short paragraph explaining what representation/basis you used for each of the calculations above and why you chose to use that representation/basis.
- (h) In the above calculations, you should have found some of the quantities to be time dependent and others to be time independent. Briefly explain why this is so. That is, for a time dependent state like  $|\Psi\rangle$  explain what makes some observables time dependent and others time independent.

5. Attached, you will find a table showing different representations of physical quantities associated with a particle-in-a-box. Make a similar table for a particle confined to a ring. Include all of the following information.

- Hamiltonian
- Eigenvalues of Hamiltonian
- Normalized eigenstates of Hamiltonian
- Coefficient of the  $n$ th eigenstate
- Probability of measuring  $E_n$
- Expectation value of Hamiltonian
- Z-component of angular momentum
- Eigenvalues of z-component of angular momentum
- Eigenstates of z-component of angular momentum
- Coefficient of  $m$ th state of z-component of angular momentum
- Probability of measuring  $m\hbar$  for z-component of angular momentum
- Expectation value of z-component of angular momentum

## Particle in a Box

	Ket Representation	Wave Function Representation	Matrix Representation
Hamiltonian	$\hat{H}$	$-\frac{\hbar^2}{2m} \frac{d^2}{dx^2}$	$\begin{pmatrix} E_1 & 0 & 0 & \dots \\ 0 & E_2 & 0 & \dots \\ 0 & 0 & E_3 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$
Eigenvalues of Hamiltonian	$E_n = \frac{\pi^2 \hbar^2}{2mL^2} n^2$	$E_n = \frac{\pi^2 \hbar^2}{2mL^2} n^2$	$E_n = \frac{\pi^2 \hbar^2}{2mL^2} n^2$
Normalized Eigenstates of Hamiltonian	$ n\rangle$	$\psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right)$	$\begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \dots$
Coefficient of $n^{\text{th}}$ energy eigenstate	$c_n = \langle n   \psi \rangle$	$c_n = \int_0^L \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right) \psi(x) dx$	$(0 \dots 1 \dots) \begin{pmatrix} c_1 \\ \vdots \\ c_n \\ \vdots \end{pmatrix}$
Probability of measuring $E_n$	$P(E_n) =  c_n ^2 =  \langle n   \psi \rangle ^2$	$P(E_n) =  c_n ^2 = \left  \int_0^L \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}x\right) \psi(x) dx \right ^2$	$P(E_n) =  c_n ^2 = \left  (0 \dots 1 \dots) \begin{pmatrix} c_1 \\ \vdots \\ c_n \\ \vdots \end{pmatrix} \right ^2$
Expectation value of Hamiltonian	$\langle \psi   H   \psi \rangle = \sum_n  c_n ^2 E_n$	$\langle \psi   H   \psi \rangle = \int_0^L \psi^*(x) \hat{H} \psi(x) dx$	$\langle \psi   H   \psi \rangle = (c_1^* \ c_2^* \ \dots) \begin{pmatrix} E_1 & 0 & \dots \\ 0 & E_2 & \dots \\ \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ \vdots \end{pmatrix}$

## Particle on a Ring

	Ket Representation	Wave Function Representation	Matrix Representation
Hamiltonian			
Eigenvalues of Hamiltonian			
Normalized Eigenstates of Hamiltonian			
Coefficient of $m^{\text{th}}$ energy eigenstates			
Probability of measuring $E_m$			
Expectation value of Hamiltonian			

## Particle on a Ring

Operator for z-component of angular momentum			
Eigenvalues of z-component of angular momentum			
Normalized Eigenstates of z-component of angular momentum			
Coefficient of $m^{\text{th}}$ eigenstates of z-component of angular momentum			
Probability of measuring $m\hbar$ for z-component of angular momentum			
Expectation value of z-component of angular momentum			