

# MTH 351 Fall 2006 – HW 5, Prob 2.

---

Due: Before class November 29

---

Consider the  $n \times n$  matrix  $A$  from Section 6.2, Problem 21 given by:

$$A = \begin{bmatrix} 1 & -1 & 0 & \cdots & 0 \\ -1 & 2 & -1 & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & -1 & 2 & -1 \\ 0 & \cdots & 0 & -1 & 2 \end{bmatrix}$$

Note that this matrix is the result of discretizing the following *two-point boundary value problem* (See Section 8.8):

$$y''(x) = f(x), \quad y'(a) = 0, \quad y(b) = 0, \quad a \leq x \leq b.$$

1. Compute (“by hand”, i.e., show each step) the LU factorization for  $A$  using
  - (a)  $n = 2$
  - (b)  $n = 3$
2. Find the pattern in the above factorizations, what do you suppose is the LU factorization for arbitrary  $n$ ?
3. Verify that the  $i, i$  entry of  $LU$  is  $A_{i,i} = 2$ .
4. Verify that the  $i, i + 1$  entry of  $LU$  is  $A_{i,i+1} = -1$ . (Since  $A$  is symmetric, we do not need to verify  $A_{i,i-1}$ .)
5. Comment on the number of floating point operations (multiplications and divisions only.. assume, rather crudely, that additions and subtractions are free!) required to solve  $Ax = b$  for this particular  $A$

(a) if given

$$A^{-1} = \begin{bmatrix} n & n-1 & n-2 & n-3 & \cdots & 1 \\ n-1 & n-1 & n-2 & n-3 & \cdots & 1 \\ n-2 & n-2 & n-2 & n-3 & \cdots & 1 \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ 1 & 1 & 1 & 1 & \cdots & 1 \end{bmatrix}$$

(b) if given  $LU$  from 2.