

DECOMPOSABLE FORMS

Let V be a vector space. A p -vector $\beta \in \wedge^p V$ is called *decomposable* if and only if there exist vectors $\alpha_i \in V$ with

$$\beta = \alpha_1 \wedge \dots \wedge \alpha_p$$

An example of an *indecomposable* 2-vector is $\alpha \wedge \beta + \gamma \wedge \delta$ where $\alpha, \beta, \gamma, \delta \in V$ are linearly independent (so that $\dim V \geq 4$).

1. Let \vec{u} be an ordinary vector in \mathbb{R}^3 , so that

$$\vec{u} = A \hat{i} + B \hat{j} + C \hat{k}$$

for some constants A, B, C . Find two vectors \vec{v} and \vec{w} such that

$$\vec{u} = \vec{v} \times \vec{w}$$

It is possible to solve this problem by brute force; find a better way if you can.

HINT: What properties should \vec{v} and \vec{w} have?

2. Show that if $\dim V = 3$ then all 2-vectors are decomposable. In other words, show that if $\{\rho, \sigma, \tau\}$ is a basis of V then

$$\gamma = A \sigma \wedge \tau + B \tau \wedge \rho + C \rho \wedge \sigma$$

is decomposable for any constants A, B, C .

HINT: Adapt your answer to the previous problem!

EXTRA CREDIT:

- Show that if $\dim V = 4$ then all 3-vectors are decomposable.
- Can you argue that all $(n-1)$ -vectors in dimension n are decomposable?