

## 1. DECOMPOSABLE FORMS

Denote the  $p$ -forms in  $\mathbb{R}^n$  by  $\wedge^p(\mathbb{R}^n)$ . A typical 1-form in  $\mathbb{R}^2$  would therefore take the form  $F = F_x dx + F_y dy \in \wedge^1(\mathbb{R}^2)$ .

A  $p$ -form  $\gamma \in \wedge^p(\mathbb{R}^n)$  is called *decomposable* if there exist 1-forms  $\alpha_i \in \wedge^1(\mathbb{R}^n)$  with

$$\gamma = \alpha_1 \wedge \dots \wedge \alpha_p$$

- (a) Show that all elements of  $\wedge^2(\mathbb{R}^3)$ , that is, all 2-forms in  $\mathbb{R}^3$ , are decomposable. In other words, show that

$$H = H_x dy \wedge dz + H_y dz \wedge dx + H_z dx \wedge dy$$

is decomposable.

*HINT: Consider the previous assignment!*

**You may cite your solution to the previous assignment without proof, so long as an explicit reference is given (“see HW #1”). If you do this, it wouldn’t hurt to include a copy of your previous assignment.**

- (b) Find an example of an *indecomposable*  $p$ -form.

*HINT: Don’t work in  $\mathbb{R}^3$ ...*

- (c) Is  $\gamma \wedge \gamma = 0$ ? Should it be? Can it be?

## 2. PICTURES OF FORMS

Let  $\alpha = 3 dx$  and  $\beta = 4 dy$ .

- (a) Draw a single picture showing the “stacks” corresponding to both  $\alpha$  and  $\beta$ .

*You may want to use different colors for the stacks corresponding to  $\alpha$  and  $\beta$ . Your drawing should be correctly scaled.*

- (b) Draw a separate picture showing the stack corresponding to  $\gamma = \alpha + \beta$ .

- (c) Choose a vector  $\vec{v} \in \mathbb{R}^2$  that is *not* parallel to the coordinate axes. Add  $\vec{v}$  to your previous diagrams and use them to compute  $\alpha(\vec{v})$ ,  $\beta(\vec{v})$ , and  $\gamma(\vec{v})$ .

*Your computation should be geometric, not algebraic.*

- (d) Did you obtain  $\gamma(\vec{v}) = \alpha(\vec{v}) + \beta(\vec{v})$ ? Should you have?