Calculus Meets Physics: 25 years of interdisciplinary discussion

Tevian Dray

Department of Mathematics Oregon State University http://www.math.oregonstate.edu/~tevian



The Paradigms in Physics Project

- Complete redesign of physics major 20 new courses
- Junior-year "paradigms" designed around common themes.
- Senior-year "capstones" finish traditional disciplinary content.
- 25+ years of continuous NSF funding.
- Living curriculum: Monthly curriculum meetings for 25+ years!
- Paradigms 2.0 implemented in 2017.
- Active engagement: 300+ documented activities!

https://paradigms.oregonstate.edu











Mathematics vs. Physics


Suppose the temperature on a rectangular slab of metal is given by

$$T(x,y) = k(x^2 + y^2)$$

where k is a constant. What is $T(r, \theta)$?


A:
$$T(r,\theta) = kr^2$$


B:
$$T(r, \theta) = k(r^2 + \theta^2)$$



Mathematics and Physics are two disciplines separated by a common language!

Geometric Reasoning





- Which vector field is conservative?
- Which vector field has nonzero curl?
- Which vector field has nonzero divergence?


Which vector field could represent an electric field? a magnetic field?

The Hill

Suppose you are standing on a hill. You have a topographic map, which uses rectangular coordinates (x,y) measured in miles. Your global positioning system says your present location is at one of the points shown. Your guidebook tells you that the height h of the hill in feet above sea level is given by

$$h = a - bx^2 - cy^2$$


where a = 5000 ft, $b = 30 \frac{\text{ft}}{\text{mi}^2}$, and $c = 10 \frac{\text{ft}}{\text{mi}^2}$.



Kinesthetic Activity

Stand up and close your eyes. Hold out your right arm in the direction of the gradient where you are standing.





Surfaces





(Each surface is dry-erasable, as are the matching contour maps.)

Raising Calculus to the Surface (Aaron Wangberg)

Raising Physics to the Surface (+ Liz Gire, Robyn Wangberg)

https://raisingcalculus.winona.edu

Thermodynamics

State Variables:

$$T = \text{temperature}$$


 $S = \text{entropy}$
 $p = \text{pressure}$
 $V = \text{volume}$

$$(U = internal energy)$$

$$dU = T dS - p dV$$

- Compressibility = $-\frac{1}{V}\frac{\partial V}{\partial p}$
- Design an experiment to measure compressibility.

Name the Experiment



$$\bullet \ \left(\frac{\partial p}{\partial T}\right)_{V} \neq \left(\frac{\partial p}{\partial T}\right)_{S}.$$

- What are the independent variables??
- Can not hold "everthing else" fixed!
- ← What is this material??

Isothermal:
$$-\frac{1}{V} \left(\frac{\partial V}{\partial p} \right)_T$$


Isoentropic: $-\frac{1}{V} \left(\frac{\partial V}{\partial p} \right)_S$

Partial Derivative Machine


- Developed for junior-level thermodynamics course
- Two positions, x_i , two string tensions (masses), F_i .
- "Find $\frac{\partial x}{\partial F}$."
- Idea: Measure Δx , ΔF ; divide.
- Mathematicians:

"That's not a derivative!"

Roundy et al., Experts' Understanding of Partial Derivatives Using the Partial Derivative Machine, PERC 2014



Thick Derivatives



Math: ∃ "bright line" between *average* rate of change and *instantaneous* rate of change.

(Such averages are used to approximate derivatives.)

Physics: "Average" refers to secant lines, not (good) approximations to tangent lines.


Move the bright line!

Thick Derivatives!

(Derivatives are fundamentally ratios of small changes, not limits.)

[Dray, AMS Blog on Education, 5/31/16]


Learning Progression



- Successively more sophisticated ways of thinking about a topic.
- Sequences supported by research on learner's ideas and skills.
- Lower anchor grounded in students' prior ideas and skills.
- *Upper anchor* grounded in knowledge and practices of experts.

Duschle et al., NRC, 2007; Plummer, 2012; Sikorski et al., 2009, 2010 Manogue, Dray, Emigh, Gire, & Roundy, PERC 2017

Multiple Representations



Extended Theoretical Framework for Concept of Derivative

Process-	Graphical	Verbal	Symbolic	Numerical	Physical
object	Slope	Rate of	Difference	Ratio of	Measurement
layer	-	Change	Quotient	Changes	
Ratio		"avg. rate of	$\frac{f(x+\Delta x)-f(x)}{\Delta x}$		T → T
		change"		numerically	
	\	"inst.	lim ···	with	T. T
Limit		rate of change"	$\Delta x \rightarrow 0$	Δx small	\rightarrow
Function		"at any point/time"	f'(x) =		tedious
				depends	repeti-
		point/ time	•••	on x	tion

No entry for symbolic differentiation!!

Roundy, Dray, Manogue, Wagner, & Weber, CRUME 18 Proceedings, MAA, 2015. https://sigmaa.maa.org/rume/Site/Proceedings.html

Differentials

Does
$$\frac{df}{dx}$$
 mean "f'(x)" or "df over dx"?
$$d(u^2) = 2u du$$

$$d(\sin u) = \cos u du$$

Instead of:

- chain rule
- related rates
- implicit differentiation
- derivatives of inverse functions
- difficulties of interpretation (units!)

One coherent idea:


"Zap equations with d"

(infinitesimal reasoning)

Dray & Manogue, CMJ 34, 283-290 (2003); CMJ 41, 90-100 (2010).

Vector Calculus

Vector calculus is about one coherent concept: Infinitesimal Displacement



$$ds = |d\vec{\mathbf{r}}|$$

$$d\vec{\mathbf{A}} = d\vec{\mathbf{r}}_1 \times d\vec{\mathbf{r}}_2$$

$$dA = |d\vec{\mathbf{r}}_1 \times d\vec{\mathbf{r}}_2|$$

$$dV = (d\vec{\mathbf{r}}_1 \times d\vec{\mathbf{r}}_2) \cdot d\vec{\mathbf{r}}_3$$

SUMMARY

- Use multiple representations, including geometry, measurement, numerical data;
- Always ask both "With respect to what," and "With what held constant."



https://math.oregonstate.edu/bridge https://books.physics.oregonstate.edu/GVC https://paradigms.oregonstate.edu https://raisingcalculus.winona.edu